arXiv:1211.5381v1 [cond-mat.stat-mech] 22 Nov 2012

Topological Rényi Entropy After a Quantum Quench

Gábor B. Halász[1,2] and Alioscia Hamma[1,3]
[1] *Perimeter Institute for Theoretical Physics, 31 Caroline Street North, Waterloo, Ontario N2L 2Y5, Canada*
[2] *Theoretical Physics, Oxford University, 1 Keble Road, Oxford OX1 3NP, United Kingdom*
[3] *Center for Quantum Information, Institute for Interdisciplinary Information Sciences, Tsinghua University, Beijing 100084, P. R. China*

We present an analytical study on the resilience of topological order after a quantum quench. The system is initially prepared in the ground state of the Toric Code Model, and then quenched by switching on an external magnetic field. During the subsequent time evolution, the variation in topological order is detected via the topological Rényi entropy of order 2. We consider two different quenches: the first one has an exact solution, while the second one requires perturbation theory. In both cases, we find that the long-term time average of the topological Rényi entropy in the thermodynamic limit is the same as its initial value. Based on our results, we argue that topological order is resilient against a wide range of quenches.

Introduction.—Topologically ordered phases in quantum many-body systems are of extreme importance in condensed matter physics [1] and in quantum information [2, 3]. They are novel phases of matter that defy the Landau paradigm of spontaneous symmetry breaking and possess a robust ground state degeneracy that makes them good candidates for quantum memories. Moreover, they feature anyonic excitations whose interactions are of a topological nature and that are therefore less subject to decoherence [2, 3]. Since these phases cannot be described by local order parameters, their characterization must be done using global properties of the ground state wavefunction. Recently, it has been argued that topological order in the ground state is revealed by an entanglement pattern called topological entropy: a universal correction to the boundary law for the entanglement entropy [4–7]. It has also been shown that this correction is robust against perturbations therefore it serves as an effective non-local order parameter for topologically ordered phases [8, 9].

Unfortunately, the entanglement entropy is extremely hard to compute and its measurement requires a complete state tomography of a possibly extended system [10]. On the other hand, there is now an increasing understanding of the fact that the Rényi entropy of order 2 contains an appreciable amount of information about the universal properties of a quantum many-body system [11]. In particular, the aforementioned topological pattern of entanglement appears in the Rényi entropy as well [6, 12]. Moreover, this quantity is significantly easier to compute and can in principle be measured directly [13]. Indeed, there are already proposals for the experimental realizations of such measurements [14].

One of the most interesting questions regarding topological order is its behavior away from equilibrium. In general, the non-equilibrium properties of quantum many-body systems are among the most fruitful subjects in both theoretical [15] and experimental [16] condensed matter physics, and they can be conveniently studied in the setting of the quantum quench: a sudden change in the parameters of the system Hamiltonian [17]. In particular, the quantum quench of a topologically ordered system was numerically studied in Ref. [18], where they found that topological order is resilient against certain types of quenches. The main disadvantage of their method is that it is only applicable to small system sizes.

In this Letter, we analytically study the behavior of a topologically ordered system after a quantum quench by determining the time evolution of the topological Rényi entropy of order 2. In particular, we prepare the system in the ground state of the Toric Code Model (TCM) [2] and quench it by switching on an external magnetic field. Concentrating on two different versions of the quench, we establish both an exact treatment and a perturbation theory. Due to the nature of these methods, our results are not confined to the small system sizes accessible by exact diagonalization [8, 18].

General formalism.—We consider the TCM with an external magnetic field in the $+z$ direction. In this model, there are $2N^2$ spins on the edges of an $N \times N$ square lattice with periodic boundary conditions [2]. The spins on the horizontal (h) and the vertical (v) edges experience different magnetic fields λ and $\kappa\lambda$, therefore the Hamiltonian takes the form [12]

$$\hat{H}(\lambda) = -\sum_{s} \hat{A}_s - \sum_{p} \hat{B}_p - \lambda \sum_{i \in h} \hat{\sigma}_i^z - \kappa\lambda \sum_{i \in v} \hat{\sigma}_i^z, \quad (1)$$

where the star operators $\hat{A}_s \equiv \prod_{i \in s} \hat{\sigma}_i^x$ and the plaquette operators $\hat{B}_p \equiv \prod_{i \in p} \hat{\sigma}_i^z$ belong to stars (s) and plaquettes (p) on the lattice containing four spins each (see Fig. 1).



FIG. 1: (Color online) Illustration of the square lattice with the physical spins located at the horizontal (black circles) and the vertical (white circles) edges. Examples of a star (red cross labeled s) and a plaquette (blue square labeled p) are included.

In the case of $\lambda = 0$, the system is exactly solvable because $[\hat{A}_s, \hat{B}_p] = [\hat{A}_s, \hat{H}(0)] = [\hat{B}_p, \hat{H}(0)] = 0$. The sectors with different sets of expectation values $\{A_s \equiv \langle \hat{A}_s \rangle = \pm 1\}$ and $\{B_p \equiv \langle \hat{B}_p \rangle = \pm 1\}$ can be treated independently. In the ground-state sector with $A_s = +1$ ($\forall s$) and $B_p = +1$ ($\forall p$), there are four linearly independent degenerate ground states that are distinguished by the topological quantum numbers $Z_{1,2} = \pm 1$. These quantum numbers are expectation values for products of $\hat{\sigma}_i^z$ operators along horizontal and vertical strings going round the lattice. The ground state with $Z_1 = Z_2 = +1$ becomes $|0\rangle = \mathcal{N} \prod_s (1 + \hat{A}_s))| \Uparrow \rangle$, where $\mathcal{N}$ is a normalization constant and $| \Uparrow \rangle$ is the completely polarized state with $\sigma_i^z \equiv \langle \hat{\sigma}_i^z \rangle = +1$ ($\forall i$).

In the case of $\lambda > 0$, the system is not exactly solvable in general because $[\hat{A}_s, \hat{\sigma}_i^z] \neq 0$. On the other hand, it is true that $[\hat{B}_p, \hat{Z}_{1,2}] = [\hat{B}_p, \hat{H}(\lambda)] = [\hat{Z}_{1,2}, \hat{H}(\lambda)] = 0$, therefore the sectors with different values of B_p and $Z_{1,2}$ can be treated independently. In the following, we only consider the lowest-energy sector with $B_p = +1$ ($\forall p$) and $Z_1 = Z_2 = +1$. The states within this sector are distinguished by the expectation values $\{A_s = \pm 1\}$, and we introduce a corresponding representation with quasi-spins A_s located at the stars [19]. In this quasi-spin representation, the operator $\hat{A}_s^z \equiv \hat{A}_s$ measures and the operator $\hat{A}_s^x$ flips the quasi-spin A_s. Up to an irrelevant additive constant, the Hamiltonian in Eq. (1) becomes

$$\hat{H}(\lambda) = -\sum_s \hat{A}_s^z - \lambda \sum_{\langle s,s' \rangle \in h} \hat{A}_s^x \hat{A}_{s'}^x - \kappa \lambda \sum_{\langle s,s' \rangle \in v} \hat{A}_s^x \hat{A}_{s'}^x, \tag{2}$$

where $\langle s, s' \rangle$ means that the summation is over edges between neighboring stars s and s'. The TCM with external magnetic field is therefore equivalent to a 2D transverse field Ising model (TFIM) in which the coupling strengths on the horizontal and the vertical edges are not equal in general.

When studying the quantum quench, we are interested in the time evolution of the quantum state $|\Psi(t)\rangle$ after a sudden change in the Hamiltonian. At time $t = 0$, the system is set up in the ground state of the initial Hamiltonian $\hat{H}(0)$ such that $|\Psi(0)\rangle = |0\rangle$. At time $t > 0$, the system is evolved with the quench Hamiltonian $\hat{H}(\lambda)$ and the state takes the general form $|\Psi(t)\rangle = \exp[-it\hat{H}(\lambda)]|0\rangle$. To extract any valuable information from this expression, we need to obtain a full solution of the Hamiltonian $\hat{H}(\lambda)$. In the following, we consider two important limits. When $\kappa = 1$, the horizontal and the vertical coupling strengths are equal, and the equivalent TFIM is the standard 2D TFIM. When $\kappa \to 0+$, the vertical coupling strength vanishes, and the equivalent TFIM factorizes into N independent 1D TFIM copies along the horizontal chains of the lattice [20]. The $\kappa = 0$ case has an exact solution available for all values of λ, while the $\kappa = 1$ case requires perturbation theory around the exactly solvable point at $\lambda = 0$.

Topological Rényi entropy.—The Rényi entropy of order α is a generalization of the usual (von Neumann) entanglement entropy that characterizes the quantum entanglement between two complementary subsystems A and $B \equiv \overline{A}$. It is defined by $S_\alpha^{AB} \equiv \log_2 \mathrm{Tr}[\hat{\rho}_A^\alpha]/(1-\alpha)$, where $\hat{\rho}_A$ is the reduced density operator for A. Note that the usual entanglement entropy is recovered in the special case of $\alpha = 1$.

The topological Rényi entropy is extracted from independent Rényi entropies $S_\alpha^{(m)}$ calculated in the four cases (m) of Fig. 2 as $S_\alpha^T \equiv -S_\alpha^{(1)} + S_\alpha^{(2)} + S_\alpha^{(3)} - S_\alpha^{(4)}$. When defining S_α^T in the standard way [5], we take $S_\alpha^{(m)} = S_\alpha^{AB}$ between the two thick subsystems A and B. However, it was argued in Ref. [12] that the presence of the $\mathbb{Z}_2$ lattice gauge structure due to the constraint $B_p = +1$ ($\forall p$) makes it possible to substitute the subsystem A with its boundary ∂A. In the following, we use the corresponding modified definition for S_α^T in which we take $S_\alpha^{(m)} = S_\alpha^{\partial A, \overline{\partial A}} \equiv S_\alpha^{\partial A}$ between the thin boundary subsystem ∂A and the rest of the system. The topological Rényi entropy is non-zero if and only if the given state has topological order. For example, its value is $S_\alpha^T = 2$ for the TCM ground state $|0\rangle$ and $S_\alpha^T = 0$ for the completely polarized state $| \Uparrow \rangle$. Note that the dimensions D and d of the subsystems need to be macroscopic such that $D > d \gg 1$.



FIG. 2: Illustration of the subsystems A and B in the four cases (m) that are used to calculate the topological Rényi entropy.

The modified definition for the topological Rényi entropy is an immense simplification to our calculations because the reduced density matrix ρ_A is then diagonal in the basis of the physical spins σ_i^z. It was shown in Ref. [12] that if the boundary subsystem ∂A consists of n closed loops with a combined length L, the Rényi entropy of order 2 for a generic state $|\Psi\rangle$ satisfying the gauge constraint $B_p = +1$ ($\forall p$) is

$$\begin{aligned} S_2^{\partial A} \; &= \; (L-n) - \log_2 \Bigg[1 + \sum_{s_1, s_2} \langle \Psi | \hat{A}_{s_1}^x \hat{A}_{s_2}^x | \Psi \rangle^2 \\ &\quad + \sum_{s_1, s_2, s_3, s_4} \langle \Psi | \hat{A}_{s_1}^x \hat{A}_{s_2}^x \hat{A}_{s_3}^x \hat{A}_{s_4}^x | \Psi \rangle^2 \; + \ldots \Bigg], \end{aligned} \tag{3}$$

where the sum contains all 2^{L-n} possible products with an even number of quasi-spin operators $\hat{A}_s^x$ chosen from each closed loop of the subsystem ∂A.

Exact treatment.—In the case of $\kappa = 0$, there are no interactions between the horizontal chains of the lattice, and the 2D system factorizes into N independent 1D systems in terms of the quasi-spins A_s. Indeed, the 2D Hamiltonian in Eq. (2) becomes the direct sum of N identical 1D Hamiltonians. If we consider any horizontal chain and label its N stars with $1 \leq l \leq N$, the corresponding 1D Hamiltonian is

$$\hat{h}(\lambda) = -\sum_{l=1}^{N} \left(\hat{A}_l^z + \lambda \hat{A}_l^x \hat{A}_{l-1}^x \right), \tag{4}$$

where the periodic boundary conditions are taken into account by $A_0 \equiv A_N$. On the other hand, the 1D TFIM Hamiltonian in Eq. (4) can be solved exactly by following a standard procedure [21]. In the first step, we map the quasi-spins A_l to fermions via the Jordan$-$Wigner transformation given by $\hat{A}_l^z = 1 - 2c_l^\dagger c_l$ and $\hat{A}_l^- = (\hat{A}_l^+)^\dagger = c_l^\dagger e^{i\pi \sum_{j=1}^{l-1} c_j^\dagger c_j}$, where $c_l^\dagger$ and c_l are standard fermionic creation and annihilation operators. The translational symmetry is then exploited by the Fourier transform $c_l = N^{-1/2} \sum_k e^{ikl} c_k$, where the sum is over the momenta $k_m = \pi(2m-1)/N$. We finally introduce new fermionic operators via the Bogoliubov transformation $c_k = \cos\theta_k \gamma_k + i \sin\theta_k \gamma_{-k}^\dagger$, and the Hamiltonian in Eq. (4) takes the free-fermion form $\hat{h}(\lambda) = \sum_k \Lambda_k (2\gamma_k^\dagger \gamma_k - 1)$, where $\gamma_k^\dagger$ and γ_k correspond to independent fermionic quasi-particles. The energies of these quasi-particles are proportional to $\Lambda_k = \sqrt{\epsilon_k^2 + \lambda^2 \sin^2 k}$ with $\epsilon_k \equiv 1 - \lambda \cos k$, and the mixing angle appearing in the Bogoliubov transformation is $\theta_k = \tan^{-1}[\lambda \sin k / (\epsilon_k + \Lambda_k)]$.

The 2D state $|\Psi(t)\rangle$ during the time evolution is the direct product of N identical 1D states $|\psi(t)\rangle$. The 1D initial state $|\psi(0)\rangle$ before the quantum quench is the 1D ground state $|\mathrm{o}\rangle$ of the initial Hamiltonian $\hat{h}(0)$ that is defined by $c_k |\mathrm{o}\rangle = 0$ $(\forall k)$. When this state is expressed in terms of the 1D ground state $|\omega\rangle$ of the quench Hamiltonian $\hat{h}(\lambda)$ that is defined by $\gamma_k |\omega\rangle = 0$ $(\forall k)$, it takes the product form $|\psi(0)\rangle = |\mathrm{o}\rangle = \prod_{k>0} (\cos\theta_k - i \sin\theta_k \gamma_k^\dagger \gamma_{-k}^\dagger) |\omega\rangle$. Note that the connections between the respective 2D and 1D ground states are $|0\rangle = \bigotimes^N |\mathrm{o}\rangle$ and $|\Omega\rangle = \bigotimes^N |\omega\rangle$. The state $|\psi(t)\rangle$ after the time evolution with $\hat{h}(\lambda)$ is then obtained by the substitution $\gamma_{\pm k}^\dagger \to \gamma_{\pm k}^\dagger e^{-2i\Lambda_k t}$ in the expression for $|\psi(0)\rangle$, and the corresponding two-operator expectation values in the position representation are given by

$$\begin{aligned}
\langle c_l c_{l'}^\dagger \rangle &\equiv \langle \psi(t) | c_l c_{l'}^\dagger | \psi(t) \rangle = \delta_{ll'} - \langle c_l^\dagger c_{l'} \rangle, \qquad (5) \\
\langle c_l^\dagger c_{l'} \rangle &= \frac{4}{N} \sum_k e^{-ik(l-l')} \sin^2\theta_k \cos^2\theta_k \sin^2(2\Lambda_k t), \\
\langle c_l^\dagger c_{l'}^\dagger \rangle &= -\langle c_l c_{l'} \rangle^* = \frac{i}{N} \sum_k e^{-ik(l-l')} \sin\theta_k \cos\theta_k \\
&\times \left[\sin^2\theta_k \left(1 - e^{-4i\Lambda_k t}\right) - \cos^2\theta_k \left(1 - e^{4i\Lambda_k t}\right) \right].
\end{aligned}$$

To calculate the Rényi entropy, we need to evaluate the quasi-spin expectation values in Eq. (3). Note that despite the factorization into independent 1D chains in terms of the quasi-spins, this calculation gives the Rényi entropy of a strongly entangled 2D state $|\Psi(t)\rangle$ in terms of the physical spins. The 2D expectation values $\langle \Psi(t) | \hat{A}_{s_1}^x \hat{A}_{s_2}^x \dots \hat{A}_{s_{2r-1}}^x \hat{A}_{s_{2r}}^x | \Psi(t) \rangle$ are simple products of independent 1D expectation values $\langle \psi(t) | \hat{A}_{l_1}^x \hat{A}_{l_2}^x \dots \hat{A}_{l_{2r-1}}^x \hat{A}_{l_{2r}}^x | \psi(t) \rangle$, where we can write $\hat{A}_l^x \hat{A}_{l'}^x = (c_l^\dagger + c_l) \prod_{j=l}^{l'-1} (1 - 2c_j^\dagger c_j)(c_{l'}^\dagger + c_{l'})$ for each pair of $\hat{A}_l^x$ operators. The resulting multi-operator expectation values can then be reduced to the two-operator expectation values in Eq. (5) by using Wick's theorem [22].



FIG. 3: (Color online) Exact long-term time average of S_2^T as a function of $\beta = \lambda/(1+\lambda)$. Three system sizes are plotted: $N = 40$ and $D = 3d = 6$ (dotted line); $N = 60$ and $D = 3d = 9$ (dashed line); $N = 80$ and $D = 3d = 12$ (solid line).

The long-term time average of the topological Rényi entropy against the magnetic field λ of the quench Hamiltonian $\hat{H}(\lambda)$ is plotted in Fig. 3. The time averages are calculated from 1000 independent time instants, which are taken from a sufficiently long interval at sufficiently large times such that $t \gg \max\{1, \lambda^{-1}\}$ for all time instants t. We notice that $\langle S_2^T \rangle$ is close to the initial value of $S_2^T = 2$ for all magnetic fields λ, and it gets closer to it if we increase the system size N as well as the subsystem dimensions D and d. We therefore argue that $S_2^T = 2$ in the thermodynamic limit for all times t after a quench with an arbitrary magnetic field λ.

Perturbation theory.—In the case of $\kappa = 1$, the Hamiltonian in Eq. (2) requires perturbation theory around the exactly solvable point at $\lambda = 0$. Our aim is to determine the time evolution of the first order correction to S_2^T. In general, the most straightforward approach is the method of perturbative continuous unitary transformations (PCUT) [12, 19, 23]. On the other hand, a naive first order perturbation theory is sufficient in our case, and we choose to derive our results using this simpler approach. Note that the same results can also be obtained from a PCUT calculation.

The unperturbed ground state of the initial Hamiltonian $\hat{H}(0)$ is $|0\rangle$ and the perturbed ground state of the quench Hamiltonian $\hat{H}(\lambda)$ is $|\Omega\rangle$ as above. In the first order of perturbation theory, the perturbed ground state can be expanded as $|\Omega\rangle = |0\rangle + (\lambda/4) \sum_{\langle \mathbf{r_1}, \mathbf{r_2} \rangle} |\{\mathbf{r_1}, \mathbf{r_2}\}\rangle$, where $|\{\mathbf{r_1}, \mathbf{r_2}\}\rangle$ is the state with two star excitations at the positions $\mathbf{r_1}$ and $\mathbf{r_2}$ such that $A_{\mathbf{r_1}}^z = A_{\mathbf{r_2}}^z = -1$. The positions of the stars are labeled by the 2D vectors $\mathbf{r} = (x, y)$ with $0 \le x, y < N$. In terms of the ground state $|\Omega\rangle$, the initial state $|0\rangle$ becomes

$$|\Psi(0)\rangle = |0\rangle = |\Omega\rangle - \frac{\lambda}{4} \sum_{\langle \mathbf{r_1}, \mathbf{r_2} \rangle} |\{\mathbf{r_1}, \mathbf{r_2}\}\rangle. \qquad (6)$$

Although the non-degenerate corrections to the excited states $|\{\mathbf{r_1}, \mathbf{r_2}\}\rangle$ are negligible in the first order calculation, the perturbation introduces a hopping between these degenerate excited states. The two star excitations can hop between neighboring stars with an amplitude $-\lambda$, and their only interaction is a hardcore repulsion: they are not allowed to be at the same star. However, since the two excitations are far away from each other in the thermodynamic limit, it is reasonable to neglect their interaction. Under this assumption, the exact eigenstates of the quench Hamiltonian with two star excitations are $|\mathbf{k_1}, \mathbf{k_2}\rangle = N^{-2} \sum_{\mathbf{r_1}, \mathbf{r_2}} e^{i(\mathbf{K_1} \cdot \mathbf{r_1} + \mathbf{K_2} \cdot \mathbf{r_2})} |\{\mathbf{r_1}, \mathbf{r_2}\}\rangle$, where $\mathbf{k} = (k_x, k_y)$ with $0 \le k_x, k_y < N$, and each excitation has a 2D momentum $\mathbf{K} = 2\pi\mathbf{k}/N$. Note that the state $|\{\mathbf{r_1},\mathbf{r_2}\}\rangle = |\{\mathbf{r_2},\mathbf{r_1}\}\rangle$ appears twice in the sum. Since the system and the initial state are both invariant under translations, we only need to consider the eigenstates with zero total momentum $\mathbf{k_1}+\mathbf{k_2}$. These states labeled by $|\mathbf{k}\rangle \equiv |\mathbf{k},-\mathbf{k}\rangle$ have relative energies $E(\mathbf{k}) = 4 - 4\lambda\epsilon(\mathbf{k})$ with respect to the ground state $|\Omega\rangle$, where $\epsilon(\mathbf{k}) \equiv \cos(K_x) + \cos(K_y)$. In terms of the eigenstates, the initial state $|\Psi(0)\rangle$ takes the form $|\Psi(0)\rangle = |0\rangle = |\Omega\rangle - (\lambda/2)\sum_{\mathbf{k}}' \epsilon(\mathbf{k})|\mathbf{k}\rangle$, where the prime means that the summation is only over half of the N^2 possible $\mathbf{k}$ values. The state $|\Psi(t)\rangle$ after the time evolution with $\hat{H}(\lambda)$ is then obtained by the substitution $\epsilon(\mathbf{k}) \to \epsilon(\mathbf{k})e^{-itE(\mathbf{k})}$, and in terms of the states $|\{\mathbf{r_1},\mathbf{r_2}\}\rangle$ it reads

$$|\Psi(t)\rangle = |\Omega\rangle + \sum_{(\mathbf{r_1},\mathbf{r_2})} Q(\mathbf{r_1},\mathbf{r_2},t)|\{\mathbf{r_1},\mathbf{r_2}\}\rangle, \tag{7}$$

where $(\mathbf{r_1},\mathbf{r_2})$ means that the summation is over pairs of stars without double counting. The coefficient of each state is $Q(\mathbf{r_1},\mathbf{r_2},t) = -(\lambda/N^2)\sum_{\mathbf{k}}' \epsilon(\mathbf{k})\cos(\mathbf{K}\cdot(\mathbf{r_1}-\mathbf{r_2}))e^{-itE(\mathbf{k})}$. Since the functions $Q(\mathbf{r_1},\mathbf{r_2},t)$ consist of many incoherent oscillations with different frequencies $E(\mathbf{k})$, the average modulus square of any such function is given by

$$\left\langle |Q(\mathbf{r_1},\mathbf{r_2},t)|^2 \right\rangle = \frac{\lambda^2}{N^4}\sum_{\mathbf{k}}{}' \epsilon(\mathbf{k})^2 \cos^2\left(\mathbf{K}\cdot(\mathbf{r_1}-\mathbf{r_2})\right) = \frac{\lambda^2}{4N^2}. \tag{8}$$

This result has a simple physical interpretation. The two excitations are always at neighboring stars before the time evolution, and the sum of the norm squares in the states with two excitations is $2N^2\lambda^2/16 = N^2\lambda^2/8$ because there are $2N^2$ ways of choosing two neighboring stars. During the time evolution, the excitations hop between stars, and the norm square is distributed uniformly between all possible states with two star excitations. Since there are $N^2(N^2-1)/2 \approx N^4/2$ ways of choosing any two stars, the uniform norm square corresponds to the result $\lambda^2/4N^2$ in Eq. (8).

When calculating the Rényi entropy for the state $|\Psi(t)\rangle$ in Eq. (7), there are two corrections to the value $S_2^{\partial A} = L - n$ for the unperturbed ground state $|0\rangle$: a static correction from the perturbed ground state $|\Omega\rangle$ and a dynamic correction from the oscillating terms $\propto Q(\mathbf{r_1},\mathbf{r_2},t)$. The static correction is linearly proportional to the combined length L of the boundary in each case (m), therefore its topological contribution vanishes [12]. On the other hand, the terms $\propto Q(\mathbf{r_1},\mathbf{r_2},t)$ give an expectation value $\langle\Psi(t)|\hat{A}^x_{\mathbf{r_1}}\hat{A}^x_{\mathbf{r_2}}|\Psi(t)\rangle = 2\,\mathrm{Re}[Q(\mathbf{r_1},\mathbf{r_2},t)]$ for each pair of stars $\mathbf{r_1}$ and $\mathbf{r_2}$ on the same closed loop of the boundary. Since $\langle \mathrm{Re}[Q(\mathbf{r_1},\mathbf{r_2},t)]^2\rangle = \langle|Q(\mathbf{r_1},\mathbf{r_2},t)|^2\rangle/2$ and there are $\ell(\ell-1)/2$ ways of choosing two stars from a closed loop of length ℓ, the time average of the dynamic correction is $\langle\Delta S_2^{\partial A}\rangle = -\lambda^2\ell(\ell-1)/(4N^2\ln 2)$ for each closed loop of the boundary in each case (m). Since there is a term $\propto \ell^2$, the topological contribution does not vanish. In particular, when the subsystem dimensions satisfy $D = 3d$, the time average of the topological Rényi entropy becomes

$$\langle S_2^T\rangle = 2 - \frac{56\lambda^2 d^2}{N^2\ln 2}. \tag{9}$$



FIG. 4: (Color online) Long-term time average of S_2^T from the exact treatment (solid line) and from the first order perturbation theory (dashed line). Two system sizes are plotted: $N = 40$ and $D = 3d = 6$ (a); $N = 80$ and $D = 3d = 12$ (b).

If we take both the system size N and the subsystem dimension d to infinity such that $d \sim \sqrt{N}$, the perturbative correction vanishes as $1/N$. We can thus choose macroscopic subsystems in the thermodynamic limit in such a way that the topological Rényi entropy is $S_2^T = 2$ for all times t.

To verify that the first order perturbation theory is indeed a good approximation at $\lambda \ll 1$ and it is not invalidated by higher order corrections, we also establish an analogous perturbation theory in the exactly solvable $\kappa = 0$ case. When $D = 3d$, the time average of the topological Rényi entropy after the first order correction takes the form

$$\langle S_2^T\rangle = 2 - \frac{\lambda^2\left(d^2+12d-1\right)}{2N\ln 2}. \tag{10}$$

The exact results and those obtained from the perturbation theory are plotted together in Fig. 4. We notice that the respective curves are in good agreement when $\lambda \ll 1$, and therefore we argue that the first order perturbation theory is a good approximation in the $\kappa = 1$ case as well.

Conclusions.—In this Letter, we studied the resilience of topological order in the TCM ground state after a quantum quench with an external magnetic field. The time evolution of topological order was detected via the topological Rényi entropy of order 2. We considered two different quenches: the integrable one was solved exactly, while the non-integrable one was treated with perturbation theory. In both cases, we found that topological order is resilient in the thermodynamic limit. Our results are in agreement with those in Ref. [18], but they apply to significantly larger system sizes.

It is interesting to discuss the generic conditions under which topological order is resilient. First of all, the results for the non-integrable quench show that integrability does not play a role. On the other hand, both quenches preserve the $\mathbb{Z}_2$ gauge structure of the TCM ground state. This gauge structure allows us to use the modified definition for S_2^T, which in turn makes the subsequent calculations possible. We therefore believe that topological order characterized by S_2^T is a robust non-equilibrium property of the system whenever gauge invariance is preserved. In perspective, it would be important to study the behavior of topological order after a more generic non-gauge-preserving quench. This further step could help us better understand the universal non-equilibrium and thermal properties of topologically ordered systems.

Acknowledgments.—We thank X.-G. Wen for illuminating discussions. This work was supported in part by the National Basic Research Program of China Grant (2011CBA00300, 2011CBA00301) and the National Natural Science Foundation of China Grant (61073174, 61033001, 61061130540). Research at the Perimeter Institute for Theoretical Physics is supported in part by the Government of Canada through NSERC and by the Province of Ontario through MRI.

[1] X.-G. Wen, *Quantum Field Theory of Many-body Systems* (Oxford University Press, 2004).
[2] A. Y. Kitaev, Ann. Phys. (N. Y.) **303**, 2 (2003).
[3] M. H. Freedman, A. Kitaev, and Z. Wang, Commun. Math. Phys. **227**, 587 (2002); C. Nayak, S. H. Simon, A. Stern, M. Freedman, and S. Das Sarma, Rev. Mod. Phys. **80**, 1083 (2008).
[4] A. Hamma, R. Ionicioiu, and P. Zanardi, Phys. Lett. A **337**, 22 (2005); A. Hamma, R. Ionicioiu, and P. Zanardi, Phys. Rev. A **71**, 022315 (2005).
[5] A. Kitaev and J. Preskill, Phys. Rev. Lett. **96**, 110404 (2006); M. Levin and X.-G. Wen, *ibid.* **96**, 110405 (2006).
[6] S. T. Flammia, A. Hamma, T. L. Hughes, and X.-G. Wen, Phys. Rev. Lett. **103**, 261601 (2009).
[7] I. H. Kim, arXiv:1210.2360.
[8] A. Hamma, W. Zhang, S. Haas, and D. A. Lidar, Phys. Rev. B **77**, 155111 (2008).
[9] S. V. Isakov, M. B. Hastings, and R. G. Melko, Nature Phys. **7**, 772 (2011).
[10] L. Amico, R. Fazio, A. Osterloh, and V. Vedral, Rev. Mod. Phys. **80**, 517 (2008).
[11] P. Zanardi and L. C. Venuti, arXiv:1205.2507.
[12] G. B. Halász and A. Hamma, arXiv:1206.4223.
[13] P. Horodecki and A. Ekert, Phys. Rev. Lett. **89**, 127902 (2002); S. J. van Enk and C. W. J. Beenakker, *ibid.* **108**, 110503 (2012).
[14] D. A. Abanin and E. Demler, Phys. Rev. Lett. **109**, 020504 (2012).
[15] A. Polkovnikov, K. Sengupta, A. Silva, and M. Vengalattore, Rev. Mod. Phys. **83**, 863 (2011).
[16] M. Greiner, O. Mandel, T. Esslinger, T. W. Hänsch, and I. Bloch, Nature **415**, 39 (2002); M. Greiner, O. Mandel, T. W. Hänsch, and I. Bloch, Nature **419**, 51 (2002).
[17] P. Calabrese and J. Cardy, Phys. Rev. Lett. **96**, 136801 (2006); M. A. Cazalilla, *ibid.* **97**, 156403 (2006); M. Rigol, V. Dunjko, V. Yurovsky, and M. Olshanii, *ibid.* **98**, 050405 (2007).
[18] D. I. Tsomokos, A. Hamma, W. Zhang, S. Haas, and R. Fazio, Phys. Rev. A **80**, 060302(R) (2009).
[19] S. Dusuel, M. Kamfor, K. P. Schmidt, R. Thomale, and J. Vidal, Phys. Rev. B **81**, 064412 (2010).
[20] J. Yu, S.-P. Kou, and X.-G. Wen, Europhys. Lett. **84**, 17004 (2008).
[21] E. Barouch, B. M. McCoy, and M. Dresden, Phys. Rev. A **2**, 1075 (1970).
[22] E. Barouch and B. M. McCoy, Phys. Rev. A **3**, 786 (1971).
[23] F. Wegner, Ann. Phys. (Leipzig) **3**, 77 (1994); S. D. Głazek and K. G. Wilson, Phys. Rev. D **48**, 5863 (1993); S. D. Głazek and K. G. Wilson, *ibid.* **49**, 4214 (1994); S. Dusuel and J. Vidal, Phys. Rev. B **71**, 224420 (2005); S. Dusuel, M. Kamfor, R. Orús, K. P. Schmidt, and J. Vidal, Phys. Rev. Lett. **106**, 107203 (2011).